


JETS IN HEAVY ION COLLISIONS.

C.A. Salgado

CERN, Theory Division, CH-1211 Geneva, Switzerland



High energetic particles traversing a dense medium lose a sizable part of their energy in form of gluon radiation. As a result, the rate of high- p_t particles is expected to be suppressed in heavy ion collisions with respect to the proton case. Recent experimental data from RHIC strongly support this scenario. This allows to study the properties of the medium by the amount of *jet quenching* it produces. The angular dependence of the radiation is modified in the medium in a characteristic way. This provides another tool to study the medium properties in a more differential measurement.

The evolution, in the vacuum, of a quark or gluon produced with high virtuality is given by gluon radiation and described by DGLAP-like evolution equations. In this way, the study of jets (production, shapes, etc...) in proton-proton, lepton-proton or e^+e^- collisions, give very valuable information about the dynamics of QCD. In heavy ion collisions, where a dense state is expected to be formed, this high energetic quark or gluon would traverse the produced medium before hadronization. In this case, the interaction with the medium modifies the gluon radiation, and hence the evolution. This has been proposed¹ long ago as a tool to study the properties of the produced medium. One of the main predictions is the suppression of particles produced at high- p_t in heavy ion collisions. Recent experimental data from RHIC found a strong suppression of the particle spectrum at large- p_t in central AuAu collisions². This same spectrum is not suppressed but enhanced in dAu collisions, also measured in RHIC³. The conclusion is that the suppression in AuAu is not due to initial state effects but to strong final state interactions.

The modified gluon radiation due to multiple (coherent) scatterings has been computed by several groups, in the multiple soft scattering limit⁴, in the hard scattering approximation⁵, they both are limiting cases of a general path integral formalism⁶. (For an alternative formalism, based on a twist-expansion see Ref. ⁷). Once the medium-modified gluon radiation is known, the modified evolution can, in principle, be computed. In Fig. 1, the $k_\perp$ -integrated spectrum




Figure 1: Medium-induced gluon energy distribution $\omega dI/d\omega$ for different media⁸. The result for $R \rightarrow \infty$ coincides with the BDMPS⁴ case.

of gluons radiated by a high energetic quark is plotted for different media in the multiple soft scattering approximation⁸ as a function of the energy of the emitted gluon ω . The quantities which characterize the medium are the length L and the transport coefficient $\hat{q}$. This last quantity is given by the amount of transverse momentum that the parton gets per unit length when suffering multiple scattering with the medium⁸. The spectrum can be written as a function of only two variables, $\omega_c = \hat{q}L^2/2$ and $R = \omega_c L$.

The first observation is that the energy distributions from Fig. 1 are cut-off at small values of the energy of the emitted gluon ω . The reason is the following, the typical transverse momentum of gluons emitted coherently is $k_\perp^2 \sim \hat{q}L$. The kinematical constraint for the $k_\perp$ -integration, $k_\perp \leq \omega$ is, then, relevant for values of $\omega/\omega_c \lesssim \sqrt{2/R}$. The positions of the maxima in the figure agrees with this estimate. One sees also that the BDMPS case⁴ is recovered⁸ for $R \rightarrow \infty$. This corresponds to remove the upper limit in the $k_\perp$ -integration.

The spectrum of gluons radiated outside a cone of angle Θ is given by^{6,8} $dI/d\omega - dI^{R'}/d\omega$, where $R' = R \cos \Theta$. This allows to compute, for a medium, the additional energy radiated outside a cone, $\Delta E(\Theta) = \int d\omega \omega (dI/d\omega - dI^{R'}/d\omega)$. In Fig. 2 this energy is plotted as a function of the angle. The induced gluon emission has a characteristic angular structure. This structure translates into modification of jet shapes due to the medium, the so called *jet broadening*.

One of the most important experimental results from RHIC is the suppression of high- p_t particles in nucleus-nucleus collisions². The recent experimental data on dAu collisions³, found an enhancement of particle production at $2 \lesssim p_t \lesssim 10$. This indicates that the suppression in AA collisions is not of initial state origin. The final state high- p_t suppression due to energy loss of the partons traveling through the produced medium is the most natural explanation for this suppression.

An simple way of computing the effect of energy loss in heavy ion collisions is through the so called quenching factor⁹

$$Q(p_t) = \frac{d\sigma^{MED}(p_t)/dp_t^2}{d\sigma^{VAC}(p_t)/dp_t^2} = \int d\epsilon P_E(\epsilon) \left(\frac{d\sigma^{VAC}(p_t(1-\epsilon))/dp_t^2}{d\sigma^{VAC}(p_t)/dp_t^2} \right), \quad (1)$$

where the medium spectrum is computed by a shift, ϵp_t , in the vacuum spectrum, $d\sigma^{VAC}$,




Figure 2: The average energy loss radiated outside an angle Θ for a quark jet of energy $E=100$ GeV.

and weighted by the probability, $P_E(\epsilon)$, that the hard parton loses a fraction ϵ of its initial energy E . The *quenching weight*, $P_E(\epsilon)$, is usually computed in the independent gluon emission approximation⁹

$$P_E(\epsilon) = \sum_{n=0}^{\infty} \frac{1}{n!} \left[\prod_{i=1}^n \int d\omega_i \frac{dI(\omega_i)}{d\omega} \right] \delta \left(\epsilon - \sum_{i=1}^n \frac{\omega_i}{E} \right) \exp \left[- \int d\omega \frac{dI}{d\omega} \right]. \quad (2)$$

Computing the suppression by means of medium-modified fragmentation functions gives very similar results⁸, so, expression (1) gives a good estimation of the effect. In Fig. 3 we compare the quenching factors that we obtain for $\omega_c=25$ GeV and $R=1000$ with the experimental data from PHENIX Collaboration² both for quark and gluon jets. For the vacuum spectrum we use the fit provided by PHENIX,

$$\frac{d\sigma^{VAC}}{dp_t^2} = \text{const} \cdot (1.71 + p_t[\text{GeV}])^{-12.44}. \quad (3)$$

The spectrum of Fig. 1 is computed in the approximation $\omega \ll E$, E being the energy of the hard parton. This is a common feature with other multiple scattering approaches^{4,5}. In the kinematical range covered by RHIC experiments, $p_t \lesssim 10$ GeV, finite energy effects could appear. In order to check the sensitivity of our results to the region $x \equiv \omega/E \sim 1$ we impose the kinematical cut for the quenching weights $P_E(x) = 0$ for $x > 1$ ^a. The energy of the initial parton is not equal to p_t ; in our estimation, we assume that a pion takes in average a fraction ~ 0.7 (~ 0.55) of the initial quark (gluon) energy¹¹. The effect of these finite energy cuts is plotted in Fig. 3. One can see that these kinematical cuts have some importance for the smallest values of p_t , however, they almost disappear for $p_t \gtrsim 10$ GeV. This conclusion depends on the actual value of ω_c ; the larger the value is, the larger the finite energy effects are too. An important observation, however, is that finite energy corrections reduce the degree of jet quenching for the

^aThe finite energy corrections to the emission vertex are the same as in the vacuum¹⁰. So, these finite energy corrections could be taken into account by simply multiplying the spectrum by the corresponding splitting function for quarks or gluons. In the case of the quarks, we have checked that the results are similar to the ones we obtain here. However, in the case of gluon jets, the configuration is symmetric and the probability of large longitudinal momentum transfer is large for $x \rightarrow 1$. Including this effect is not simple in the present formalism, as the hadronization of emitted gluons is not included.




Figure 3: Quenching factors for quarks and gluons compared with PHENIX data² on π^0 suppression in AuAu collisions. Solid lines are computed using the spectrum of Fig. 1 for $R=1000$, $\omega_c=25$ GeV. Dashed lines take into account finite energy cuts. Dotted line takes into account the p_t -dependence of the fractional contribution from quarks and gluons to the final pions.

small values of p_t . This changes the curvature of the suppression in agreement with the trend of experimental data. The contribution of quarks and gluons to the observed pion yield depends on p_t . For comparison, we have computed the suppression taking into account the fractional contribution of quarks and gluons to the final particle from Ref. ¹². This gives us an estimation of the net effect and the slopes. A more refined analysis, including nuclear parton distribution functions, fragmentation function, etc.. is, however, needed for a quantitative description.

1. M. Gyulassy and M. Plumer, Phys. Lett. B **243** (1990) 432; M. Gyulassy and X. n. Wang, Nucl. Phys. B **420** (1994) 583.
2. B. B. Back *et al.* [PHOBOS Collaboration], arXiv:nucl-ex/0302015.
J. Adams *et al.* [STAR Collaboration], arXiv:nucl-ex/0305015.
S. S. Adler *et al.* [PHENIX Collaboration], arXiv:nucl-ex/0304022.
3. S. S. Adler *et al.* [PHENIX Collaboration], arXiv:nucl-ex/0306021.
J. Adams *et al.* [STAR Collaboration], arXiv:nucl-ex/0306024.
B. B. Back [PHOBOS Collaboration], arXiv:nucl-ex/0306025.
I. Arsene [BRAHMS Collaboration], arXiv:nucl-ex/0307003.
4. R. Baier, Y. L. Dokshitzer, A. H. Mueller, S. Peigne and D. Schiff, Nucl. Phys. B **483** (1997) 291; Nucl. Phys. B **484** (1997) 265. B. G. Zakharov, JETP Lett. **65** (1997) 615.
5. M. Gyulassy, P. Levai and I. Vitev, Phys. Rev. Lett. **85** (2000) 5535; Nucl. Phys. B **594** (2001) 371.
6. U. A. Wiedemann, Nucl. Phys. B **588** (2000) 303; Nucl. Phys. A **690** (2001) 731.
7. X.f. Guo and X.N. Wang, Phys.Rev.Lett. **85** (2000) 3591; Nucl.Phys. A **696** (2001) 788.
8. C. A. Salgado and U. A. Wiedemann, Phys. Rev. Lett. **89** (2002) 092303; Nucl. Phys. A **715** (2003) 783; arXiv:hep-ph/0302184.
9. R. Baier, Y. L. Dokshitzer, A. H. Mueller and D. Schiff, JHEP **0109** (2001) 033.
10. R. Baier, Y. L. Dokshitzer, A. H. Mueller and D. Schiff, Nucl. Phys. B **531** (1998) 403.
11. K. J. Eskola and H. Honkanen, Nucl. Phys. A **713** (2003) 167.

12. S. R. Klein and R. Vogt, Phys. Rev. C **67** (2003) 047901.